Table 5.12d Adjusted Patient Survival, <u>Living Donor</u> Kidney Transplants Survival at 3 Months, 1 Year, 5 Years, and 10 Years

	3	Months		1 Year		5	5 Years		10 Years				
	(Tx 2006 - 2007)			(Tx 2	(Tx 2006 - 2007)			(Tx 2002 - 2007)			(Tx 1997 - 2007)		
	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.	
Total													
All	11,177	99.6%	0.1%	11,177	98.7%	0.1%	34,288	91.9%	0.2%	56,493	77.3%	0.3%	
Age at Tx													
<1 Year	6	99.8%	52.4%	6	99.8%	52.4%	15	86.6%	8.6%	35	94.2%	3.9%	
1-5 Years	157	100.0%	0.0%	157	100.0%	3.4%	544	96.7%	0.8%	960	95.2%	0.8%	
6-11 Years	127	+	+	127	+	+	490	98.0%	0.7%	957	94.8%	1.1%	
12-17 Years	279	100.0%	0.0%	279	100.0%	1.9%	1,084	97.3%	0.6%	2,001	91.4%	1.0%	
18-34 Years	2,119	99.8%	0.1%	2,119	99.2%	0.2%	6,844	95.5%	0.3%	12,416	87.8%	0.5%	
35-49 Years	3,325	99.7%	0.1%	3,325	99.1%	0.2%	10,312	93.7%	0.3%	17,601	81.7%	0.5%	
50-64 Years	3,948	99.3%	0.1%	3,948	98.2%	0.2%	11,621	89.2%	0.4%	17,890	67.8%	0.7%	
65+ Years	1,216	99.2%	0.3%	1,216	96.9%	0.5%	3,378	79.2%	1.0%	4,633	43.6%	2.0%	
Recipient Ethnicity/Race													
White	7,457	99.5%	0.1%	7,457	98.6%	0.1%	22,812	91.5%	0.3%	37,954	76.6%	0.4%	
African American	1,632	99.7%	0.1%	1,632	98.7%	0.3%	5,238	90.6%	0.5%	8,567	73.4%	0.9%	
Hispanic/Latino	1,495	99.6%	0.2%	1,495	99.1%	0.2%	4,473	94.5%	0.5%	7,152	83.1%	0.9%	
Asian	460	99.8%	0.2%	460	99.4%	0.3%	1,376	95.2%	0.8%	2,239	84.1%	1.7%	
Other/Multi-race	133	100.0%	0.0%	133	99.3%	0.6%	386	89.1%	2.1%	578	80.4%	2.9%	
Unknown	0	-	-	0	-	-	3	57.8%	22.0%	3	57.4%	22.0%	
Recipient Gender													
Female	4,268	99.6%	0.1%	4,268	98.9%	0.2%	13,632	92.1%	0.3%	22,868	77.4%	0.5%	
Male	6,909	99.5%	0.1%	6,909	98.6%	0.1%	20,656	91.7%	0.3%	33,625	77.2%	0.4%	

Source: OPTN/SRTR Data as of May 4, 2009.

Notes

Survival rates are adjusted to the characteristics of the 3 month and 1 year cohort. See Technical Notes for details.

^{(+) =} Values not determined due to insufficient follow-up. (-) = Values not determined since there were no transplants in the category.

Cohorts are transplants performed during the years listed at the top of the table.

Patient survival follows patients from first transplant of this type until death. Counts for patient and graft survival are different because a patient may have more than one transplant for a type of organ.

Center volume = Center's yearly transplants performed during the base period, based on kidney and kidney-pancreas transplants. Multi-organ transplants are excluded.

Table 5.12d (Continued) Adjusted Patient Survival, Living Donor Kidney Transplants Survival at 3 Months, 1 Year, 5 Years, and 10 Years

	3	Months			1 Year			5 Years			10 Years		
	(Tx 2006 - 2007)			(Tx 2006 - 2007)			(Tx 2002 - 2007)			(Tx 1997 - 2007)			
	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.	
Total All	11,177	99.6%	0.1%	11,177	98.7%	0.1%	34,288	91.9%	0.2%	56,493	77.3%	0.3%	
Primary Diagnosis													
Glomerular Diseases	3,151	99.7%	0.1%	3,151	99.1%	0.2%	9,886	94.8%	0.3%	16,936	84.7%	0.5%	
Diabetes Hypertensive	2,436	99.5%	0.1%	2,436	98.5%	0.2%	7,498	86.1%		12,196	61.6%	0.9%	
Nephrosclerosis	1,918	99.4%	0.2%	1,918	98.3%	0.3%	5,141	90.0%	0.6%	7,691	72.9%	1.09	
Polycystic Kidneys Tubular and Interstitial	1,253	99.7%	0.1%	1,253	99.5%	0.2%	3,715	96.4%	0.4%	5,649	87.1%	0.99	
Diseases Renovascular & Other	613	99.4%	0.3%	613	98.6%	0.4%	2,126	90.1%	0.9%	3,814	78.1%	1.29	
Vascular Diseases Congenital, Rare Familial, & Metabolic	235	99.3%	0.5%	235	98.2%	0.8%	1,072	90.8%	1.1%	1,989	70.7%	1.99	
Disorders	377	99.6%	0.4%	377	98.0%	0.9%	1,329	94.7%	1.0%	2,258	83.3%	1.99	
Neoplasms	43	+	+	43	+	+	144	88.8%	3.1%	240	71.0%	6.49	
Other	1,132	99.7%	0.2%	1,132	98.3%	0.4%	3,324	91.7%	0.7%	5,646	78.9%	1.19	
Unknown	19	100.0%	0.0%	19	96.1%	3.8%	53	79.7%	8.0%	74	65.4%	10.19	

Source: OPTN/SRTR Data as of May 4, 2009.

Notes:

Multi-organ transplants are excluded.

^{(+) =} Values not determined due to insufficient follow-up. (-) = Values not determined since there were no transplants in the category.

Cohorts are transplants performed during the years listed at the top of the table.

Patient survival follows patients from first transplant of this type until death. Counts for patient and graft survival are different because a patient may have more than one transplant for a type of organ.

Center volume = Center's yearly transplants performed during the base period, based on kidney and kidney-pancreas transplants.

Survival rates are adjusted to the characteristics of the 3 month and 1 year cohort. See Technical Notes for details.

Table 5.12d (Continued) Adjusted Patient Survival, Living Donor Kidney Transplants Survival at 3 Months, 1 Year, 5 Years, and 10 Years

	3 Months				1 Year		Ę	5 Years		10 Years			
	(Tx 2	2006 - 20	007)	(Tx 2	2006 - 20	007)	(Tx 2	2002 - 20	007)	(Tx 1997 - 2007)			
	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.	
Total All	11,177	99.6%	0.1%	11,177	98.7%	0.1%	34,288	91.9%	0.2%	56,493	77.3%	0.3%	
Transplant Center State													
Alabama	280	+	+	280	+	+	896	90.7%	1.4%	1,446	75.1%	2.1%	
Arkansas	48	100.0%	0.0%	48	98.2%	1.7%	192	91.1%	2.5%	389	69.9%	3.8%	
Arizona	304	99.7%	0.3%	304	99.2%	0.4%	811	92.2%	1.2%	1,198	77.8%	2.7%	
California	1,202	99.8%	0.1%	1,202	99.0%	0.3%	3,724	93.2%	0.6%	6,256	79.6%	1.0%	
Colorado	198	99.6%	0.4%	198	98.7%	0.7%	510	92.8%	1.5%	841	76.4%	3.09	
Connecticut	107	100.0%	0.0%	107	99.2%	0.8%	297	92.6%	2.0%	623	78.9%	2.7%	
District of Columbia	157	+	+	157	+	+	518	93.9%	1.3%	954	73.8%	2.8%	
Delaware	9	+	+	9	+	+	18	90.0%	8.2%	24	91.2%	7.3%	
Florida	381	99.3%	0.4%	381	98.4%	0.6%	1,119	89.4%	1.4%	1,815	74.7%	2.0%	
Georgia	216	100.0%	0.0%	216	99.2%	0.6%	661	92.6%	1.4%	1,026	79.0%	2.6%	
Hawaii	25	96.5%	3.4%	25	96.5%	3.4%	105	91.5%	3.2%	150	83.5%	4.4%	
Iowa	65	100.0%	0.0%	65	98.8%	1.2%	257	90.0%	2.4%	472	78.4%	2.9%	
Illinois	605	99.5%	0.3%	605	98.6%	0.5%	1,790	91.3%	0.9%	2,737	79.0%	1.69	
Indiana	176	100.0%	0.0%	176	99.0%	0.7%	433	96.7%	1.2%	687	84.2%	2.59	
Kansas	37	+	+	37	+	+	124	94.1%	2.8%	221	78.7%	4.89	
Kentucky	64	98.6%	1.4%	64	98.6%	1.4%	203	92.4%	2.6%	360	85.5%	3.09	
Louisiana	135	99.3%	0.7%	135	98.0%	1.1%	434	91.1%	1.8%	701	74.3%	3.19	
Massachusetts	367	99.5%	0.3%	367	99.3%	0.4%	1,067	92.4%		1,814	80.3%		
Maryland	299	99.4%	0.4%	299	97.5%	0.8%	922	92.1%	1.1%	1,820	75.3%	1.69	
Maine	47	100.0%	0.0%	47	98.2%	1.7%	139	92.7%	2.4%	235	77.4%	1	
Michigan	425	99.5%	0.3%	425	98.2%	0.6%	1,411	90.2%		2,407	75.4%	1	
Minnesota	579	99.3%	0.3%	579	98.3%	0.5%	1,843	91.7%	!	3,003	77.8%	!	
Missouri	239	99.7%	0.3%	239	99.0%	0.6%	666	94.1%	1.3%	1,198	76.8%	1	
Mississippi	3	+	+	3	+	+	8	+	+	24	73.2%	1	
North Carolina	215	99.1%	0.6%	215	97.4%	1.0%	813	89.7%	ļ	1,418	78.6%	1	
North Dakota	41	+	+	41	+	+	146	93.6%		230	76.4%	1	
Nebraska 	112	99.3%		112	99.3%	0.7%	281	95.5%	Į.	410	83.2%	1	
New Hampshire	37	+	+	37	+	+	96	95.4%	ļ	143	82.7%	!	
New Jersey	329	99.5%	0.4%	329	98.7%	0.6%	1 -	92.7%	Į.	1,537	79.6%		
New Mexico	35	+	+	35	+	+	121	89.3%		216	84.1%	1	
Nevada	22	+	+	22	+	+	83	90.4%	3.8%	164	67.3%	1	
New York	938	99.5%		938	98.7%	0.4%	2,658	93.0%		4,250	76.9%	1	
Ohio	644	99.7%	0.2%	644	98.2%	0.5%	1,954	90.1%	0.9%	2,948	71.0%	1.99	

(Continued)

Source: OPTN/SRTR Data as of May 4, 2009.

Notes:

^{(+) =} Values not determined due to insufficient follow-up. (-) = Values not determined since there were no transplants in the category.

Cohorts are transplants performed during the years listed at the top of the table.

Patient survival follows patients from first transplant of this type until death. Counts for patient and graft survival are different because a patient may have more than one transplant for a type of organ.

Center volume = Center's yearly transplants performed during the base period, based on kidney and kidney-pancreas transplants.

Multi-organ transplants are excluded.

Survival rates are adjusted to the characteristics of the 3 month and 1 year cohort. See Technical Notes for details.

Table 5.12d (Continued) Adjusted Patient Survival, <u>Living Donor</u> Kidney Transplants Survival at 3 Months, 1 Year, 5 Years, and 10 Years

	3 Months				1 Year			5 Years			10 Years		
	(Tx 2006 - 2007)			(Tx 2006 - 2007)			(Tx 2002 - 2007)			(Tx 1997 - 2007)			
	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.	
Transplant Center State													
Oklahoma	39	97.8%	2.2%	39	95.6%	3.0%	167	88.9%	2.8%	289	75.9%	4.2%	
Oregon	150	99.4%	0.5%	150	98.9%	0.8%	536	93.0%	1.4%	856	79.9%	2.8%	
Pennsylvania	565	99.4%	0.3%	565	98.8%	0.4%	1,779	92.3%	0.8%	2,745	78.0%	1.5%	
Puerto Rico	21	+	+	21	+	+	114	90.8%	3.3%	253	64.7%	5.5%	
Rhode Island	48	+	+	48	+	+	167	92.5%	2.1%	290	69.6%	5.5%	
South Carolina	63	98.5%	1.4%	63	98.5%	1.4%	157	92.4%	2.6%	329	70.8%	4.1%	
South Dakota	29	100.0%	0.0%	29	94.0%	4.1%	100	87.9%	4.0%	123	83.7%	4.7%	
Tennessee	245	99.6%	0.4%	245	97.3%	1.0%	802	89.8%	1.4%	1,309	74.5%	2.2%	
Texas	649	99.3%	0.3%	649	98.5%	0.5%	1,931	91.3%	0.8%	3,154	78.3%	1.4%	
Utah	171	100.0%	0.0%	171	99.5%	0.5%	457	94.4%	1.4%	771	83.7%	2.2%	
Virginia	327	99.5%	0.4%	327	98.9%	0.5%	1,032	89.2%	1.3%	1,632	75.2%	2.2%	
Vermont	27	97.3%	2.6%	27	94.5%	3.7%	59	95.9%	2.3%	93	85.7%	7.5%	
Washington	201	+	+	201	+	+	696	91.1%	1.4%	1,269	80.2%	1.7%	
Wisconsin	279	99.7%	0.3%	279	99.1%	0.5%	891	93.6%	1.0%	1,488	78.5%	2.0%	
West Virginia	22	+	+	22	+	+	91	90.3%	4.2%	175	67.7%	6.2%	

Source: OPTN/SRTR Data as of May 4, 2009.

Notes

(+) = Values not determined due to insufficient follow-up. (-) = Values not determined since there were no transplants in the category.

Cohorts are transplants performed during the years listed at the top of the table.

Patient survival follows patients from first transplant of this type until death. Counts for patient and graft survival are different because a patient may have more than one transplant for a type of organ.

Center volume = Center's yearly transplants performed during the base period, based on kidney and kidney-pancreas transplants.

Multi-organ transplants are excluded.

Survival rates are adjusted to the characteristics of the 3 month and 1 year cohort. See Technical Notes for details.